

Modified dispersionless Veselov–Novikov equations and corresponding hydrodynamic chains

Maxim V. Pavlov

P.N. Lebedev Physical Institute, Moscow, Russia

Abstract

Various links connecting well-known hydrodynamic chains and corresponding 2+1 nonlinear equations are described.

Contents

1	Introduction	1
2	Dispersionless Veselov–Novikov equations	2
3	Modified VN equation	4
	References	7

keywords: Riemann mapping, hydrodynamic chain, Gibbons equation

MSC: 35L40, 35L65, 37K10; PACS: 02.30.J, 11.10.E.

1 Introduction

Theory of integrable hydrodynamic chains started with famous paper [1] (see also, [2], [6], [7], [8], [9], [10], [12], [20]). Recent investigations are obtained in [3] and in [15] (see also, [5], [4]).

This paper is devoted to a description of various links associated with the integrable hydrodynamic chain

$$B_{t^1}^k = B_{t^0}^{k+1} + \frac{1}{2} B^0 B_{t^0}^k + (k+1) B^k B_{t^0}^0, \quad k \in \mathbf{Z}. \quad (1)$$

which is a particular case of extended Kupershmidt hydrodynamic chains (see [11], [16])

$$B_{t^1}^k = B_{t^0}^{k+1} + \frac{1}{\beta} B^0 B_{t^0}^k + (k+\gamma) B^k B_{t^0}^0, \quad k \in \mathbf{Z}.$$

This hydrodynamic chain can be written in the Hamiltonian form

$$B_{t^1}^k = [(2k+1) B^{k+n} D_{t^0} + (2n+1) D_{t^0} B^{k+n}] \frac{\delta \mathbf{H}_1}{\delta B^n},$$

where the Hamiltonian is given by $\mathbf{H}_1 = \int [B^1/3 + (B^0)^2/4] dt^0$. The extended Kupershmidt hydrodynamic chains possess an infinite series of conservation laws. All positive conservation law densities h_n are polynomial with respect to positive moments B^k . Negative conservation law densities h_{-n} are more complicated. For instance, $h_{-1} = -2 \exp(B^{-1}/2)$, $h_{-2} = -B^{-2} \exp(3B^{-1}/2)/3$. Thus, the above hydrodynamic chain possesses an infinite series of commuting flows. For instance, two first negative flows are

$$B_{t^{-1}}^k = e^{B^{-1}/2} [B_{t^0}^{k-1} - k B^{k-1} B_{t^0}^{-1}], \quad (2)$$

$$B_{t^{-2}}^k = e^{3B^{-1}/2} \left[B_{t^0}^{k-2} + \frac{B^{-2}}{2} B_{t^0}^{k-1} - \left(\frac{3}{2} k B^{k-1} B^{-2} + (k-1) B^{k-2} \right) B_{t^0}^{-1} - k B^{k-1} B_{t^0}^{-2} \right]$$

2 Dispersionless Veselov–Novikov equations

The first negative commuting flow (2) is invariant (see [16]) under a transformation of independent variables $t^0 \leftrightarrow t^{-1}$ and the substitution

$$\tilde{B}^k = B^{-k-2} e^{(k+1)B^{-1}}.$$

All other commuting flows are invariant under a transformation of independent variables $t^{k-1} \leftrightarrow t^{-k}$. For instance, the second commuting flow (3) transforms into (1), while (1) transforms into (3).

Let us *combine* these commuting flows, i.e. we have new integrable hydrodynamic chain

$$\begin{aligned} B_t^k &= B_{t^0}^{k+1} + \frac{B^0}{2} B_{t^0}^k + (k+1) B^k B_{t^0}^0 \\ &+ e^{3B^{-1}/2} \left[B_{t^0}^{k-2} + \frac{B^{-2}}{2} B_{t^0}^{k-1} - \left(\frac{3}{2} k B^{k-1} B^{-2} + (k-1) B^{k-2} \right) B_{t^0}^{-1} - k B^{k-1} B_{t^0}^{-2} \right] \end{aligned} \quad (4)$$

determined by the *mixed* Hamiltonian $\mathbf{H} \equiv \mathbf{H}_1 + \mathbf{H}_{-2} = \int [B^1/3 + (B^0)^2/4 - B^{-2} \exp(3B^{-1}/2)/3] dt^0$ (see (1) and (3)).

Introducing new field variables

$$v = \frac{B^0}{2}, \quad u = e^{B^{-1}/2}, \quad w = \frac{B^{-2}}{2} e^{B^{-1}},$$

two conservation laws (see (2))

$$\partial_{t^{-1}} \frac{B^0}{2} = \partial_{t^0} e^{B^{-1}/2}, \quad \partial_{t^{-1}} e^{B^{-1}/2} = \partial_{t^0} \left(\frac{B^{-2}}{2} e^{B^{-1}} \right)$$

can be written in Egorov's form (see [19], [15])

$$v_{t^{-1}} = u_{t^0}, \quad u_{t^{-1}} = w_{t^0}.$$

These conservation laws together with the conservation law (see (4))

$$u_t = (uv)_{t^0} + (uw)_{t^{-1}}$$

determine a dispersionless limit of the remarkable symmetric Veselov–Novikov equation

$$\Omega_{t^0 t^{-1} t} = (\Omega_{t^0 t^{-1}} \Omega_{t^0 t^0})_{t^0} + (\Omega_{t^0 t^{-1}} \Omega_{t^{-1} t^{-1}})_{t^{-1}}, \quad (5)$$

where

$$v = \Omega_{t^0 t^0}, \quad u = \Omega_{t^0 t^{-1}}, \quad w = \Omega_{t^{-1} t^{-1}}.$$

Another 2+1 quasilinear system of the first order can be derived from the Kupershmidt hydrodynamic chain (1) and its first higher commuting flow

$$B_{t^2}^k = B_{t^0}^{k+2} + \frac{3}{2} B^0 B_{t^0}^{k+1} + \left(\frac{B^1}{2} + \frac{5(B^0)^2}{8} \right) B_{t^0}^k + (k+2) B^{k+1} B_{t^0}^0 + (k+1) B^k \left(B_{t^0}^1 + \frac{5B^0}{2} B_{t^0}^0 \right)$$

determined by the Hamiltonian $\mathbf{H}_2 = \int [B^2/5 + B^0 B^1/2 + 5(B^0)^3/24] dt^0$.

First two conservation laws of the Kupershmidt hydrodynamic chain (1)

$$B_{t^1}^0 = \partial_{t^0} \left(B^1 + \frac{3}{4} (B^0)^2 \right), \quad \partial_{t^1} \left(B^1 + \frac{3}{4} (B^0)^2 \right) = \partial_{t^0} \left(B^2 + 2B^0 B^1 + \frac{3}{4} (B^0)^3 \right)$$

and first conservation law of its above higher commuting flow

$$B_{t^2}^0 = \partial_{t^0} \left(B^2 + \frac{5}{2} B^0 B^1 + \frac{25}{24} (B^0)^3 \right)$$

can be written together as a dispersionless limit of non-symmetric Veselov–Novikov equation

$$\Omega_{t^1 t^1} = \Omega_{t^0 t^2} - \Omega_{t^0 t^0} \Omega_{t^0 t^1} + \frac{1}{3} \Omega_{t^0 t^0}^3, \quad (6)$$

where

$$\Omega_{t^0 t^1} = \frac{B^1}{2} + \frac{3}{8} (B^0)^2, \quad \Omega_{t^0 t^2} = \frac{B^2}{2} + \frac{5}{4} B^0 B^1 + \frac{25}{48} (B^0)^3.$$

Remark: The symmetric dispersionless Veselov–Novikov equation (5) can be obtained from the compatibility condition $\partial_{t^1}(\partial_{t^{-1}} p) = \partial_{t^{-1}}(\partial_{t^1} p)$, where the generating function of conservation law densities p satisfies

$$p_{t^{-1}} = -\partial_{t^0} \frac{u}{p}, \quad p_t = \partial_{t^0} \left(\frac{p^3}{3} + vp - \frac{uw}{p} - \frac{u^3}{3p^3} \right),$$

while the non-symmetric dispersionless Veselov–Novikov equation (6) can be obtained from the compatibility condition $\partial_{t^1}(\partial_{t^2} p) = \partial_{t^2}(\partial_{t^1} p)$, where the generating function of conservation law densities p satisfies

$$p_{t^1} = \partial_{t^0} \left(\frac{p^3}{3} + \Omega_{t^0 t^0} p \right), \quad p_{t^2} = \partial_{t^0} \left[\frac{p^5}{5} + \Omega_{t^0 t^0} p^3 + (\Omega_{t^0 t^1} + \Omega_{t^0 t^0}^2) p \right].$$

Corresponding Gibbons equations (see [6], [17]) associated with the above hydrodynamic chains are

$$\begin{aligned}
\lambda_{t^1} - (p^2 + v) \lambda_{t^0} &= \frac{\partial \lambda}{\partial p} \left[p_{t^1} - \partial_{t^0} \left(\frac{p^3}{3} + vp \right) \right], \\
\lambda_{t^{-2}} - \left(\frac{uw}{p^2} + \frac{u^3}{p^4} \right) \lambda_{t^0} &= \frac{\partial \lambda}{\partial p} \left[p_{t^{-2}} + \partial_{t^0} \left(\frac{uw}{p} + \frac{u^3}{3p^3} \right) \right], \\
\lambda_t - \left(p^2 + v + \frac{uw}{p^2} + \frac{u^3}{p^4} \right) \lambda_{t^0} &= \frac{\partial \lambda}{\partial p} \left[p_t - \partial_{t^0} \left(\frac{p^3}{3} + vp - \frac{uw}{p} - \frac{u^3}{3p^3} \right) \right], \\
\lambda_{t^{-1}} - \frac{u}{p^2} \lambda_{t^0} &= \frac{\partial \lambda}{\partial p} \left(p_{t^{-1}} + \partial_{t^0} \frac{u}{p} \right), \\
\lambda_{t^2} - (p^4 + 3vp^2 + v^2 + s) \lambda_{t^0} &= \frac{\partial \lambda}{\partial p} \left[p_{t^2} - \partial_{t^0} \left(\frac{p^5}{5} + vp^3 + (v^2 + s)p \right) \right],
\end{aligned}$$

where $v = \Omega_{t^0 t^0}$, $s = \Omega_{t^0 t^1}$, and (6) can be written as 2+1 quasilinear system of the first order

$$\partial_{t^1} v = \partial_{t^0} s, \quad \partial_{t^1} s = \partial_{t^2} v + \partial_{t^0} \left(\frac{v^3}{3} - vs \right).$$

Remark: Since the function Ω is unique for both Veselov–Novikov equations, both of them possess the same set of hydrodynamic reductions determined by the Gibbons–Tsarev system (see [8], [16]).

3 Modified VN equation

The generating function of conservation laws and commuting flows (see [16]) is given by

$$\partial_{\tau(\zeta)} p(\lambda) = \frac{1}{2} \partial_{t^0} \ln \frac{p(\lambda) - p(\zeta)}{p(\lambda) + p(\zeta)}, \quad (7)$$

where $p(\lambda)$ can be found due to the Bürmann–Lagrange series (see, for instance, [13]) from the Riemann mappings

$$\lambda|_{p \rightarrow \infty} = p^2 \exp \left(\sum_{k=0}^{\infty} \frac{B^k}{p^{2(k+1)}} \right), \quad \lambda|_{p \rightarrow 0} = p^{-2} \exp \left(\sum_{k=1}^{\infty} B^{-k} p^{2(k-1)} \right).$$

In this section we consider another expansion

$$p(\lambda) = \tilde{h}_0 + \lambda \tilde{h}_1 + \lambda^2 \tilde{h}_2 + \dots, \quad \lambda \rightarrow 0, \quad (8)$$

where λ is another local parameter. In such a case, (7) leads to an infinite series of generating functions

$$\partial_{\tau^0} p(\lambda) = \frac{1}{2} \partial_{t^0} \ln \frac{p(\lambda) - \tilde{h}_0}{p(\lambda) + \tilde{h}_0}, \quad \partial_{\tau^1} p(\lambda) = -\partial_{t^0} \frac{\tilde{h}_1 p(\lambda)}{p^2(\lambda) - \tilde{h}_0^2}, \dots,$$

where

$$\partial_{\tau(\lambda)} = \partial_{\tau^0} + \lambda \partial_{\tau^1} + \lambda^2 \partial_{\tau^2} + \dots$$

Introducing a new generating function of conservation law densities

$$\tilde{p}(\lambda) = \ln \frac{p(\lambda) + \tilde{h}_0}{p(\lambda) - \tilde{h}_0}, \quad (9)$$

and eliminating t^0 , both above generating functions can be written in the form

$$\partial_{\tau^1} \tilde{p}(\lambda) = \partial_{\tau^0} \left(\frac{\tilde{h}_1}{\tilde{h}_0} \sinh \tilde{p}(\lambda) \right) \quad (10)$$

which is a first coefficient from (7) (see (9) and [15])

$$\partial_{\tau(\zeta)} \tilde{p}(\lambda) = \partial_{\tau^0} \ln \frac{e^{\tilde{p}(\lambda)} - e^{\tilde{p}(\zeta)}}{e^{\tilde{p}(\lambda) + \tilde{p}(\zeta)} - 1}.$$

Substituting the Taylor expansion (8) into the above generating functions, one can obtain new 2+1 nonlinear equations

$$\Omega_{\tau^0 \tau^0} = \frac{1}{2} \ln \frac{\Omega_{t^0 \tau^1}}{\Omega_{t^0 \tau^0}}, \quad \Omega_{\tau^1 \tau^1} = \Omega_{\tau^0 \tau^2} - \Omega_{\tau^0 \tau^1}^2 + \frac{1}{2} \exp(4\Omega_{\tau^0 \tau^0}),$$

where $\tilde{h}_k \equiv \Omega_{t^0 \tau^k}$, because $p(\lambda) \equiv \Omega_{t^0 \tau(\lambda)}$ (the last formula is a consequence of (7), if $p \rightarrow \infty$).

Remark: The second above equation was found in [15]. This is a nontrivial generalization of a continuum limit of discrete KP hierarchy (see, for instance, [14]). Thus, we described links between different well-known 2+1 integrable equations in this paper.

Theorem: *The hydrodynamic chain*

$$A_{\tau^1}^k = \frac{A_{\tau^0}^{k+1}}{A^0} - \frac{(k+2)A^{k+1} - kA^{k-1}}{(A^0)^2} A_{\tau^0}^0, \quad k = 0, 1, 2, \dots \quad (11)$$

satisfies the Gibbons equation

$$\lambda_{\tau^1} - \frac{\cosh \tilde{p}}{A^0} \cdot \lambda_{\tau^0} = \lambda_{\tilde{p}} \left(\tilde{p}_{\tau^1} - \partial_{\tau^0} \frac{\sinh \tilde{p}}{A^0} \right), \quad (12)$$

where the equation of the Riemann surface is given by

$$\lambda = \sum_{n=0}^{\infty} \frac{A^n}{\cosh^{n+1} \tilde{p}}. \quad (13)$$

Proof: The substitution (11) and (13) in (12) leads to an identity.

Theorem: *The above hydrodynamic chain (11) is Hamiltonian*

$$A_{\tau^1}^k = [(k+2)A^{k+1} - kA^{k-1}] D_{\tau^0} + A_{\tau^0}^{k+1} \left] \frac{\partial h}{\partial A^0}, \right.$$

where the Hamiltonian density $h = \ln A^0$.

Proof: Integrable hydrodynamic chains associated with more general Hamiltonian structure were considered in [18].

Remark: Simplest hydrodynamic reductions

$$a_{\tau^1}^i = \partial_{\tau^0} \frac{\sinh a^i}{A^0}$$

are given by the moment decomposition

$$A^k = \frac{1}{k+1} \sum_{i=1}^N \epsilon_i (\cosh a^i)^{k+1}, \quad k = 0, 1, 2, \dots, \quad \sum_{i=1}^N \epsilon_i = 0.$$

Theorem: 2+1 nonlinear equation

$$\Omega_{\tau^1 \tau^1} = \Omega_{\tau^0 \tau^2} - \frac{1}{2} \Omega_{\tau^0 \tau^1}^2 + \epsilon \exp(-2\Omega_{\tau^0 \tau^0})$$

can be determined from the compatibility condition $\partial_{\tau^1}(\partial_{\tau^2} p) = \partial_{\tau^2}(\partial_{\tau^1} p)$, where the generating functions of conservation laws are given by

$$p_{\tau^1} = \partial_{\tau^0} [e^{-U}(e^p + \epsilon e^{-p})], \quad p_{\tau^2} = \partial_{\tau^0} \left[\frac{e^{2p} - \epsilon^2 e^{-2p}}{2} e^{-2U} - V(e^p - \epsilon e^{-p}) e^{-U} \right],$$

where $U = \Omega_{\tau^0 \tau^0}$, $V = \Omega_{\tau^0 \tau^1}$.

Remark: Substituting the transformation

$$\mu = e^{-U}(e^p + \epsilon e^{-p}) - V$$

into the above generating functions and eliminating τ^0 , one can obtain the generating function of conservation laws

$$\mu_{\tau^2} = \partial_{\tau^1} \left[\frac{\mu + V}{2} \sqrt{(\mu + V)^2 - 4\epsilon e^{-2U}} - V(\mu + V) - W \right]$$

generalizing ($\epsilon = 0$) the generating function of conservation laws for the Benney hydrodynamic chain

$$\mu_{\tau^2} = \partial_{\tau^1} \left[\frac{\mu^2}{2} - \frac{V^2}{2} - W \right],$$

where $W = \Omega_{\tau^0 \tau^2}$. Such a transformation for $\epsilon = 0$ was found first by Yuji Kodama (see, for instance, [14]).

Let us introduce new notations $U = -C^{-1}$, $V = -C^0$ and $q = p - U$.

Theorem: The integrable hydrodynamic chain

$$C_{\tau^1}^{-1} = C_{\tau^0}^0, \quad C_{\tau^1}^0 = \partial_{\tau^0} (C^1 - \epsilon e^{-2C^{-1}}), \quad (14)$$

$$C_{\tau^1}^k = C_{\tau^0}^{k+1} + k C^k C_{\tau^0}^0 - \epsilon e^{-2C^{-1}} [C_{\tau^0}^{k-1} + 2(k-1) C^{k-1} C_{\tau^0}^{-1}], \quad k = 1, 2, \dots$$

satisfies the Gibbons equation (see (10))

$$\lambda_{\tau^1} - (e^q - \epsilon e^{-2C^{-1}-q})\lambda_{\tau^0} = \frac{\partial \lambda}{\partial q} \left[q_{\tau^1} - \partial_{\tau^0} \left(e^q + C^0 + \epsilon e^{-2C^{-1}-q} \right) \right], \quad (15)$$

where the Riemann mapping is determined by

$$\lambda = e^q + \sum_{n=0}^{\infty} C^n e^{-nq}. \quad (16)$$

Proof: Indeed, a substitution the above series and the hydrodynamic chain (14) into the Gibbons equation (15) leads to an identity.

This Riemann mapping (16) is the same as for a continuum limit of two-dimensional Toda lattice. This hydrodynamic chain (14) is a member of an integrable hierarchy generalizing ($\epsilon = 0$) an integrable hierarchy containing a continuum limit of two-dimensional Toda lattice.

Acknowledgement

I thank Leonid Bogdanov, Eugeni Ferapontov and Boris Kupershmidt for their stimulating and clarifying discussions.

I am grateful to the Institute of Mathematics in Taipei (Taiwan) where some part of this work has been done, and especially to Jen-Hsu Chang, Jyh-Hao Lee, Ming-Hien Tu and Derchyi Wu for fruitful discussions.

References

- [1] *D.J. Benney*, Some properties of long non-linear waves, *Stud. Appl. Math.*, **52** (1973) 45-50.
- [2] *B.A. Dubrovin*, Hamiltonian formalism of Whitham-type hierarchies and topological Landau-Ginsburg models, *Comm. Math. Phys.*, **145** (1992) 195-207. *B.A. Dubrovin*, Geometry of 2D topological field theories, *Lecture Notes in Math.* 1620, Springer-Verlag (1996) 120-348.
- [3] *E.V. Ferapontov, K.R. Khusnutdinova*, On integrability of (2+1)-dimensional quasi-linear systems, *Comm. Math. Phys.*, **248** (2004) 187-206, *E.V. Ferapontov, K.R. Khusnutdinova*, The characterization of 2-component (2+1)-dimensional integrable systems of hydrodynamic type, *J. Phys. A: Math. Gen.*, **37** No. 8 (2004) 2949-2963.
- [4] *E.V. Ferapontov, K.R. Khusnutdinova, M.V. Pavlov*, Classification of integrable (2+1)-dimensional quasilinear hierarchies, *Theor. Math. Phys.* **144** (2005) 35-43.
- [5] *E.V. Ferapontov, D.G. Marshall*, Differential-geometric approach to the integrability of hydrodynamic chains: the Haantjes tensor, *arXiv:nlin.SI/0505013*.

- [6] *J. Gibbons*, Collisionless Boltzmann equations and integrable moment equations, *Physica D*, **3** (1981) 503-511.
- [7] *J. Gibbons, Yu. Kodama*, Solving dispersionless Lax equations. In N. Ercolani et al., editor, *Singular limits of dispersive waves*, v. **320** of NATO ASI Series B, page 61. Plenum (1994) New York.
- [8] *J. Gibbons, S.P. Tsarev*, Reductions of the Benney equations, *Phys. Lett. A* **211** (1996) 19-24. *J. Gibbons, S.P. Tsarev*, Conformal maps and reductions of the Benney equations, *Phys. Lett. A* **258** (1999) 263-271.
- [9] *Yu. Kodama*, A method for solving the dispersionless KP equation and its exact solutions. *Phys. Lett. A*, **129** No. 4 (1988) 223-226. *Yu. Kodama*, A solution method for the dispersionless KP equation, *Prog. Theor. Phys. Supplement.* **94** (1988) 184.
- [10] *I.M. Krichever*, The averaging method for two-dimensional "integrable" equations, *Funct. Anal. Appl.* **22** No. 3 (1988) 200-213, *I.M. Krichever*, Spectral theory of two-dimensional periodic operators and its applications, *Russian Math. Surveys* **44** No. 2 (1989) 145-225. *I.M. Krichever*, The dispersionless equations and topological minimal models, *Comm. Math. Phys.*, **143** No. 2 (1992) 415-429. *I.M. Krichever*, The τ -function of the universal Whitham hierarchy, matrix models and topological field theories, *Comm. Pure Appl. Math.* **47** (1994) 437-475.
- [11] *B.A. Kupershmidt*, Deformations of integrable systems, *Proc. Roy. Irish Acad. Sect. A* **83** No. 1 (1983) 45-74. *B.A. Kupershmidt*, Normal and universal forms in integrable hydrodynamical systems, *Proceedings of the Berkeley-Ames conference on nonlinear problems in control and fluid dynamics* (Berkeley, Calif., 1983), in *Lie Groups: Hist., Frontiers and Appl. Ser. B: Systems Inform. Control, II*, Math Sci Press, Brookline, MA, (1984) 357-378.
- [12] *B.A. Kupershmidt, Yu.I. Manin*, Long wave equation with free boundaries. I. Conservation laws. *Func. Anal. Appl.*, **11** No. 3 (1977) 188-197. *B.A. Kupershmidt, Yu.I. Manin*, Long wave equations with a free surface. II. The Hamiltonian structure and the higher equations. *Func. Anal. Appl.*, **12** No. 1 (1978) 20-29. *D.R. Lebedev, Yu.I. Manin*, Conservation laws and representation of Benney's long wave equations, *Phys. Lett. A*, **74** No. 3,4 (1979) 154-156. *D.R. Lebedev*, Benney's long wave equations: Hamiltonian formalism, *Lett. Math. Phys.*, **3** (1979) 481-488.
- [13] *M.A. Lavrentiev, B.V. Shabat*, *Metody teorii funktsiï kompleksnogo peremennogo* (Russian) [Methods of the theory of functions of a complex variable] Third corrected edition Izdat. "Nauka", Moscow (1965) 716 pp. *P. Henrici*, *Metody teorii funktsiï kompleksnogo peremennogo* Topics in computational complex analysis. IV. The Lagrange-Bürmann formula for systems of formal power series. Computational aspects of complex analysis (Braunlage, 1982), 193-215, NATO Adv. Sci. Inst. Ser. C, Math. Phys. Sci., 102, Reidel, Dordrecht, 1983.
- [14] *L.A. Yu*, Waterbag reductions of the dispersionless discrete KP hierarchy, *J. Phys. A: Math. Gen.*, **33** (2000) 8127-8138.

- [15] *M.V. Pavlov*, Classification of the Egorov hydrodynamic chains, Theor. Math. Phys., **138** No. 1 (2004) 55-71.
- [16] *M.V. Pavlov*, The Kupershmidt hydrodynamic chains and lattices. *to appear in IMRN*.
- [17] *M.V. Pavlov*, The Hamiltonian approach in the classification and the integrability of hydrodynamic chains. *to appear in J. Math. Phys.*
- [18] *M.V. Pavlov*, Algebro-geometric approach in the theory of integrable hydrodynamic type systems. *to appear in Comm. Math. Phys.*
- [19] *M. V. Pavlov, S.P. Tsarev*, Three-Hamiltonian structures of the Egorov hydrodynamic type systems, Funct. Anal. Appl., **37** No. 1 (2003) 32-45.
- [20] *V.E. Zakharov*, Benney's equations and quasi-classical approximation in the inverse problem method, Funct. Anal. Appl., **14** No. 2 (1980) 89-98. *V.E. Zakharov*, On the Benney's Equations, Physica 3D (1981) 193-200.